

Differential Geometry

Thomas Brosnan

Notes taken in Professor Sergey Frolov's class, Michaelmas Term 2024

“hokay” -Sergey Frolov

Contents

| | | |
|----------|--|-----------|
| 1 | Definition of a Manifold | 5 |
| 1.1 | Regions | 5 |
| 1.2 | Differentiable Manifold | 5 |
| 1.3 | Abuse of notation | 5 |
| 2 | Elements of Topology | 6 |
| 2.1 | Topological space | 6 |
| 2.2 | Metric space | 6 |
| 2.3 | Orientation | 7 |
| 3 | Mappings on Manifolds | 8 |
| 3.1 | Manifold mappings | 8 |
| 3.2 | Equivalent manifolds | 8 |
| 3.3 | Tangent vector | 8 |
| 3.4 | Push forward | 8 |
| 3.5 | Directional derivative | 9 |
| 3.6 | Riemann metric | 9 |
| 4 | Tensors | 10 |
| 4.1 | Tensor def | 10 |
| 4.2 | Operations on Tensors | 10 |
| 4.3 | Co-Vectors | 11 |
| 4.4 | Skew-Symmetric Tensor | 11 |
| 4.5 | Volume element | 11 |
| 4.6 | Generalized push forward | 12 |
| 4.7 | Pull back | 12 |
| 5 | Manifolds and surfaces | 13 |
| 5.1 | Immersion | 13 |
| 5.2 | Manifold with boundary | 13 |
| 5.3 | Surfaces as Manifolds | 13 |
| 5.4 | Orientation of surfaces | 13 |
| 5.5 | Two-sided hyper-surface | 14 |
| 6 | Lie Groups | 15 |
| 6.1 | Group | 15 |
| 6.2 | Lie Group | 15 |
| 6.3 | Example of Lie groups | 15 |
| 7 | Projective spaces | 17 |
| 7.1 | Real protective space | 17 |
| 7.2 | Quaternions | 17 |
| 7.3 | Complex Projective spaces | 17 |
| 8 | Lie Algebras | 18 |
| 8.1 | Neighborhood of identity element | 18 |
| 8.2 | Commutator | 18 |
| 8.3 | Lie Algebra | 19 |

| | | |
|-----------|---|-----------|
| 9 | One parameter subgroups | 20 |
| 9.1 | Co-ords of the first kind | 20 |
| 9.2 | Co-ords of the second kind | 20 |
| 10 | Linear Representations | 21 |
| 10.1 | Representations | 21 |
| 10.2 | Schur's Lemma | 21 |
| 10.3 | Push Forward Representation | 21 |
| 10.4 | Faithful | 21 |
| 10.5 | Inner automorphism | 22 |
| 10.6 | One Parameter Adjoint | 22 |
| 11 | Simple Lie Algebras and Forms | 23 |
| 11.1 | Simple & Semi-Simple | 23 |
| 11.2 | Killing Form | 23 |
| 12 | Group Actions | 24 |
| 12.1 | Left and Right actions | 24 |
| 12.2 | Transitivity | 24 |
| 12.3 | Examples of Homogeneous spaces | 24 |
| 13 | Vector Bundles | 26 |
| 13.1 | Tangent Bundle | 26 |
| 13.2 | Cotangent Bundle | 26 |
| 13.3 | Symplectic Manifold | 26 |
| 14 | Vector and Tensor Fields | 27 |
| 14.1 | Vector Field | 27 |
| 14.2 | Tensor Field | 27 |
| 14.3 | Commutator or Lie Bracket | 27 |
| 14.4 | Integral curves | 28 |
| 14.5 | Exponential function of Vector Fields | 28 |
| 15 | The Lie Derivative | 29 |
| 15.1 | Action of Flows on Tensors | 29 |
| 15.2 | Lie Derivative | 29 |
| 16 | Covariant Differenciation | 30 |

1 Definition of a Manifold

1.1 Regions

- A *region* (“open set”) is a set of D points in \mathbb{R}^n such that together with each point p_0 , D also contains all points sufficiently closer to p_0 , i.e.:

$$\forall p_0 = (x_0^1, \dots, x_0^n) \in D \exists \epsilon > 0, \\ \text{st } p = (x^1, \dots, x^n) \in D, \text{ iff } |x^i - x_0^i| < \epsilon.$$

- A *region with out a boundary* is obtained from a region D by adjoining all boundary points to D . The *boundary* of a region is the set of all boundary points.

1.2 Differentiable Manifold

- A differentiable n -dimensional manifold is a set M together with the following structure on it. The set M is the union of a finite or countably infinite collection of subsets U_q with the following properties:
 - Each subset U_q has defined on it co-ords $x_q^\alpha, \alpha = 1, \dots, n$ called local co-ords by virtue of which U_q is identifiable with a region of Euclidean n -space \mathbb{R}^n with Euclidean co-ords x_q^α . The U_q with their co-ord systems are called *charts* or *local coordinate neighborhoods*.
 - Each non-empty intersection $U_q \cap U_p$ of a pair of charts thus has defined on it two co-ord systems, the restriction of x_p^α and x_q^α . It is required that under each of these coordinatizations the intersection $U_q \cap U_p$ is identifiable with a region of \mathbb{R}^n and that each of these co-ordinate systems be expressible in terms of the other in a one to one differentiable manner. Thus, if a *transition* functions from x_p^α to x_q^α and back are given by:

$$x_p^\alpha = x_p^\alpha(x_q^1, \dots, x_q^n), \quad \alpha = 1, \dots, n \\ x_q^\alpha = x_q^\alpha(x_p^1, \dots, x_p^n), \quad \alpha = 1, \dots, n$$

Then the *Jacobian* $J_{pq} = \det(\partial x_p^\alpha / \partial x_q^\alpha)$ is non-zero on $U_p \cap U_q$.

1.3 Abuse of notation

- Regular partial derivative do not have the same “canceling” that total derivative have ($dx * dy / dx = dy$) But we can restore this property through Einstein summation convention. That is that:

$$\sum_{\gamma=1}^n \frac{\partial x_p^\alpha}{\partial x_q^\gamma} \frac{\partial x_q^\gamma}{\partial x_q^\beta} = \frac{\partial x_p^\alpha}{\partial x_q^\gamma} \frac{\partial x_q^\gamma}{\partial x_q^\beta} = \delta_\beta^\alpha$$

2 Elements of Topology

2.1 Topological space

- A topological space is a set X of points of which certain subsets called *open sets* of the topological space, are distinguished, these open sets have to satisfy:
 - The intersection of any two (and hence of any finite collection) open sets should again be an open set.
 - The union of any collection of open sets must again be open.
 - The empty set and the whole set X must be open.
- The compliment of any open set is called a *closed* set of the topological space.

In Euclidean space \mathbb{R}^n the “Euclidean topology” is the usual one where the open sets are the open regions.

2.1.1 Induced topology

- Given any subset $A \in \mathbb{R}^n$, the *induced topology* on A is that where the open sets are the intersections $A \cap U$, where U ranges over all open sets of \mathbb{R}^n .

2.1.2 Continuity

- A map $f : X \rightarrow Y$ of one topological space to another is called *continuous* if the complete inverse image $f^{-1}(U)$ of every open set $U \subset Y$ is open in X .

2.1.3 Homeomorphic

- Two topological space are *topologically equivalent* or *homeomorphic* if there is a one to one and onto map (bijective) between them, such that it and its inverse are continuous.

2.1.4 Topology on a manifold

- The topology on a manifold M is given by the following specifications of the open sets. In every local co-ordinate neighborhood U_q the open regions are to be open in the topology on M ; the totality of open sets of M is then obtained by admitting as open, also arbitrary unions countable collections of such regions, i.e. by closing under countable unions.

2.2 Metric space

- A *metric space* is a set which comes equipped with a “distance function” i.e. a real-valued function $\rho(x, y)$, defined on pairs x, y of its elements and having the following properties:
 - Symmetry: $\rho(x, y) = \rho(y, x)$.
 - Positivity: $\rho(x, x) = 0$, $\rho(x, y) > 0$ if $x \neq y$.
 - The triangle inequality: $\rho(x, y) \leq \rho(x, z) + \rho(z, y)$.

2.2.1 Hausdorff

- A topological space is called *Hausdorff* if any two points are contained in disjoint open sets. Any metric space is Hausdorff because the open balls of radius $\rho(x, y)/3$ with centers at x, y do not intersect.

All topological spaces we consider will be Hausdorff.

2.2.2 Compact

- A topological space X is said to be compact if every countable collection of open sets covering X contains a finite sub-collection already covering X .

If X is a metric space the compactness is equivalent to the condition that from every sequence of points of X a convergent sub-sequence can be selected.

2.2.3 Connected

- A topological space is connected if any two points can be joined by a continuous path.

2.3 Orientation

- A manifold M is said to be *orientated* if one can choose its atlas (collection of all the charts) so that for every pair U_p, U_q of intersecting co-ordinate neighborhoods the Jacobian of the transition functions is positive.
- We say that the co-ordinate systems x and y define the *same orientation* if $J > 0$ and the *opposite orientation* if $J < 0$.

3 Mappings on Manifolds

3.1 Manifold mappings

- A mapping $f : M \rightarrow N$ is said to be smooth of smoothness class k if for all p, q for which f determines functions $y_q^b(x_p^1, \dots, x_p^m) = f(x_p^1, \dots, x_p^m)_p^b$, these functions are, where defined, smooth of smoothness class k (i.e. all their partial derivatives up to those of k -th order exist and are continuous).

the smoothness class of f cannot exceed the maximum class of the manifolds.

3.2 Equivalent manifolds

- The manifolds M and N are said to be *smoothly equivalent* or *diffeomorphic* if there is a one to one and onto map f such that both $f : M \rightarrow N$ and $f^{-1} : N \rightarrow M$ are smooth of some class $k \geq 1$.

Since f^{-1} exists then the Jacobian $J_{pq} \neq 0$ wherever it is defined.

3.3 Tangent vector

- A *tangent* vector to an m -dim manifold M at an arbitrary point x is represented in terms of local co-ords x_p^α by an m tuple ξ^α of components which are linked to the components in terms of any other system x_q^β of local co-ords by:

$$\xi_p^\alpha = \left(\frac{\partial x_p^\alpha}{\partial x_q^\beta} \right)_x \xi_q^\beta, \quad \forall \alpha \quad (3.1)$$

- The set of all tangent vectors to an m -dim manifold M at a point x forms an m -dim vector space $T_x = T_x M$, the *tangent space* to M at the point x .
- Thus, the velocity at x of any smooth curve M through x is a tangent vector to M at x .

3.4 Push forward

- A smooth map f from M to N gives rise for each x to a *push forward* or an *induced linear* map to tangent spaces:

$$f_* : T_x M \rightarrow T_{f(x)} N$$

defined as sending the velocity at x of any smooth curve $x = x(\tau)$ on M to the velocity vector at $f(x)$ of the curve $f(x(\tau))$ on N . If the map f is given by: $y^b = f^b(x^1, \dots, x^m)$ for $x \in M$ and $y \in N$, then the push forward map f_* is:

$$\xi^\alpha \rightarrow \eta^b = \frac{\partial f^b}{\partial x^\alpha} \xi^\alpha.$$

- For a real valued function $f : M \rightarrow \mathbb{R}$, the push-forward map f_* corresponding to each $x \in M$ is a real valued linear function on the tangent space to M at x :

$$\xi^\alpha \rightarrow \eta = \frac{\partial f}{\partial x^\alpha} \xi^\alpha$$

and it is represented by the gradient of f at x , and is a co-vector or one form. Thus f_* can be identified with the differential df , in particular:

$$dx_p^\alpha : \xi^\alpha \rightarrow \eta = \xi_p^\alpha$$

3.5 Directional derivative

- We can associate with each vector $\xi = (\xi^i)$ a linear differential operator as follows: Since the gradient $\frac{\partial f}{\partial x^i}$ of a function f is a co-vector, the quantity:

$$\partial_\xi f = \xi^i \frac{\partial f}{\partial x^i}$$

is a scalar called the directional derivative of f in the direction of ξ .

- Thus an arbitrary vector ξ corresponds to the operator:

$$\partial_\xi = \xi^i \frac{\partial}{\partial x^i}$$

So we can identify $\frac{\partial}{\partial x^i} \equiv e_i$ as the *Canonical basis of the tangent space*.

3.6 Riemann metric

- A *Riemann metric* on a manifold M is a point-dependent, positive-definite quadratic form on the tangent vectors at each point, depending smoothly on the local co-ords of the points.

Thus at each point $x = (x_p^1, \dots, x_p^m)$ of each chart U_p , the metric is given by a symmetric metric $g_{\alpha\beta}(x_p^1, \dots, x_p^m)$, and determines a symmetric scalar product of pairs of tangent vectors at the point x .

$$\langle \xi, \eta \rangle = g_{\alpha\beta}^{(p)} \xi_p^\alpha \eta_p^\beta = \langle \eta, \xi \rangle, \quad |\xi|^2 = \langle \xi, \xi \rangle$$

This scalar product is to be co-ordinate independent:

$$g_{\alpha\beta}^{(p)} \xi_p^\alpha \eta_p^\beta = g_{\alpha\beta}^{(q)} \xi_q^\alpha \eta_q^\beta$$

And therefor the coefficients $g_{\alpha\beta}^{(p)}$ of the quadratic form transform as:

$$g_{\gamma\delta}^{(q)} = \frac{\partial x_p^\alpha}{\partial x_q^\gamma} \frac{\partial x_p^\beta}{\partial x_q^\delta} g_{\alpha\beta}^{(p)} \quad (3.2)$$

For a *pseudo-Riemann* metric M one just requires the quadratic form to be *nondegenerate*, (i.e. the determinant of g is not 0). Note that 3.2 can be re-written as:

$$ds^2 = g_{\alpha\beta}^{(p)} dx_p^\alpha dx_p^\beta = g_{\alpha\beta}^{(q)} dx_q^\alpha dx_q^\beta$$

Where ds is called a line element, and it is chart-independent. ds is used to measure the distance between two infinitesimally close points.

4 Tensors

4.1 Tensor def

- A *tensor of type* (k, l) and rank $k + l$ on an m -dim manifold M is given each local co-ord system (x_p^i) by a family of functions:

$${}^{(p)}T_{j_1, \dots, j_l}^{i_1, \dots, i_k}(x) \text{ of the point } x.$$

In other local co-ord (x_q^i) the components ${}^{(p)}T_{j_1, \dots, j_l}^{i_1, \dots, i_k}(x)$ of the same tensor are:

$${}^{(p)}T_{t_1, \dots, t_l}^{s_1, \dots, s_k}(x) = \frac{\partial x_q^{s_1}}{\partial x_p^{i_1}} \dots \frac{\partial x_q^{s_k}}{\partial x_p^{i_k}} \frac{\partial x_p^{j_1}}{\partial x_q^{t_1}} \dots \frac{\partial x_p^{j_l}}{\partial x_q^{t_l}} \cdot {}^{(p)}T_{j_1, \dots, j_l}^{i_1, \dots, i_k}(x)$$

4.2 Operations on Tensors

4.2.1 Permutation of indices

- Let σ be some permutation of $1, 2, \dots, l$. σ acts on the ordered tuple (j_1, \dots, j_l) as $\sigma(j_1, \dots, j_l) = (j_{\sigma_1}, \dots, j_{\sigma_l})$. We say that a tensor $\tilde{T}_{j_1, \dots, j_l}^{i_1, \dots, i_k}(x)$ is obtained from a tensor $T_{j_1, \dots, j_l}^{i_1, \dots, i_k}(x)$ by means of a permutation σ of the lower indices if at each point of M :

$$\tilde{T}_{j_1, \dots, j_l}^{i_1, \dots, i_k}(x) = T_{\sigma(j_1, \dots, j_l)}^{i_1, \dots, i_k}(x)$$

Permutation of upper indicies is defined similarly.

4.2.2 Contraction of indicies

- By the contraction of a tensor $T_{j_1, \dots, j_l}^{i_1, \dots, i_k}(x)$ of type (k, l) with respect to the indicies i_a, j_a we mean the tensor (summation over n):

$$T_{j_1, \dots, j_{l-1}}^{i_1, \dots, i_{k-1}}(x) = T_{j_1, \dots, j_{a-1}, n, j_{a+1}, \dots, j_l}^{i_1, \dots, i_{a-1}, n, i_{a+1}, \dots, i_k}(x)$$

Of type $(k-1, l-1)$

4.2.3 Product of Tensors

- Given two tensors $T = (T_{j_1, \dots, j_l}^{i_1, \dots, i_k})$ of type (k, l) and $P = (P_{j_1, \dots, j_q}^{i_1, \dots, i_p})$ of type (p, q) , we define their product to be the tensor product $S = T \otimes P$ of type $(k+p, l+q)$ with components:

$$S_{j_1, \dots, j_{l+q}}^{i_1, \dots, i_{k+p}} = T_{j_1, \dots, j_l}^{i_1, \dots, i_k} P_{j_{l+1}, \dots, j_q}^{i_{k+1}, \dots, i_p}$$

This multiplication is *not commutative* but it is associative.

- The result of applying the above three operations to tensors are again tensors.

4.3 Co-Vectors

- Recall that the differential of a function f of x^1, \dots, x^n corresponding to the increments dx^i in the x^i is:

$$df = \frac{\partial f}{\partial x^i} dx^i$$

Since dx^i is a vector df has the same value in any co-ord system. In general, given any co-vector (T_i) , the differential form $T_i dx^i$ is invariant under change of chart. We can thus identify $dx^i \equiv e^i$ as the *canonical basis of co-vectors or cotangent space*.

4.4 Skew-Symmetric Tensor

- A *skew-symmetric tensor* of type $(0, k)$ is a tensor T_{i_1, \dots, i_k} satisfying:

$$T_{\sigma(i_1, \dots, i_k)} = \mathfrak{s}(\sigma) T_{i_1, \dots, i_k}$$

where for all permutations $\mathfrak{s}(\sigma)$ is the sign function. i.e. $\mathfrak{s}(\sigma) = +1(-1)$ for even(odd) permutation. If two indices of T_{i_1, \dots, i_k} are the same then the corresponding component of T_{i_1, \dots, i_k} is 0. This means if $k > n$ the tensor is automatically 0.

- The standard basis at a given point is:

$$dx^{i_1} \wedge \dots \wedge dx^{i_k}, \quad i_1 < i_2 < \dots < i_k$$

Where:

$$dx^{i_1} \wedge \dots \wedge dx^{i_k} = \sum_{\sigma \in S_k} \mathfrak{s}(\sigma) e^{i_{\sigma_1}} \otimes \dots \otimes e^{i_{\sigma_k}}$$

Here S_k is the symmetric group. i.e. the group of all permutations of k elements.

- The differential form of the skew-symmetric tensor (T_{i_1, \dots, i_k}) is:

$$\begin{aligned} T_{i_1, \dots, i_k} e^{i_1} \otimes \dots \otimes e^{i_k} &= \sum_{i_1 < i_2 < \dots < i_k} T_{i_1, \dots, i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k} \\ &= \frac{1}{k!} T_{i_1, \dots, i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k} \end{aligned}$$

Where the last step can be made as both $dx^{i_1} \wedge \dots \wedge dx^{i_k}$ and T_{i_1, \dots, i_k} are anti-symmetric.

4.5 Volume element

- A metric g_{ij} on a manifold is a tensor of type $(0, 2)$ and on an oriented manifold of $\dim(M) = n$ such a metric gives rise to a *volume element*:

$$T_{i_1, \dots, i_n} = \sqrt{|g|} \epsilon_{i_1, \dots, i_n}, \quad g = \det(g_{ij})$$

It is convenient to write the volume element in the notation of differential forms:

$$\Omega = \sqrt{|g|} dx^1 \wedge \dots \wedge dx^n$$

If g_{ij} is Riemann then the *volume* V of M is:

$$V = \int_M \Omega = \int_M \sqrt{|g|} dx^1 \wedge \dots \wedge dx^n$$

4.6 Generalized push forward

- We can generalize the push forward map we had on vectors earlier to the space of tensors $(k, 0)$:

$$f_* : \xi^{i_1, \dots, i_k} \rightarrow \eta^{a_1, \dots, a_k} = \frac{\partial f^{a_1}}{\partial x^{i_1}} \cdots \frac{\partial f^{a_k}}{\partial x^{i_k}} \xi^{i_1, \dots, i_k}$$

4.7 Pull back

- Let $T_x^{(0,k)}M$ denote the space of tensors of type $(0, k)$ at $x \in M$. Let f be a smooth map from M to N . It gives rise to a map:

$$f^* : T_{f(x)}^{(0,k)}N \rightarrow T_x^{(0,k)}M$$

which in terms of $x^i \in U \subset M$, and $y^a \in V \subset N$ is written as:

$$f^* : \eta_{a_1, \dots, a_k} \rightarrow \xi_{i_1, \dots, i_k} = \frac{\partial f^{a_1}}{\partial x^{i_1}} \cdots \frac{\partial f^{a_k}}{\partial x^{i_k}} \eta_{a_1, \dots, a_k}$$

The map f^* is called the *pullback*.

- We can then note the following relationship between pullbacks and push forwards. Let us denote the action of a vector on another vector as follows:

$$\zeta(\theta) \equiv \zeta_{i_1, \dots, i_k} \theta^{i_1, \dots, i_k}$$

Then we can write that:

$$(f^*\eta)(\xi) = \frac{\partial f^{a_1}}{\partial x^{i_1}} \cdots \frac{\partial f^{a_k}}{\partial x^{i_k}} \eta_{a_1, \dots, a_k} \xi^{i_1, \dots, i_k} = \eta(f_*\xi)$$

5 Manifolds and surfaces

5.1 Immersion

- A manifold M of dim m is said to be immersed in a manifold N of dim $n \geq m$ if \exists a smooth map $f : M \rightarrow N$ such that the push forward map f_* is at each point a one to one map of the tangent space.

The map f is called the *immersion* of M to N .

Since f_* is at each a point one to one map of the tangent space, in terms of local co-ords the Jacobian matrix of f at each point has rank equal to $m = \dim M$.

5.1.1 Embedding

- An immersion of M to N is called an *embedding* if it one to one. Then M is called a *sub-manifold* of N .
- To see the difference between these two definitions note that a Klein bottle is immersed in \mathbb{R}^3 but not embedded as its tangent spaces are distinct (intersecting points can have different tangent spaces) but the map of points is not one- to one as there are cross overs.

5.2 Manifold with boundary

- A closed region A of a manifold M defined by an inequality:

$$f(x) \leq 0, \quad (\text{or } f(x) \geq 0)$$

where f is a real-valued function on M . This region is a *Manifold with boundary*. It is assumed that the boundary ∂A given by $f(x) = 0$ is a non-singular sub-manifold of M i.e. $\nabla f \neq 0$ on ∂A .

5.2.1 Closed manifold

- A compact manifold without a boundary is called *closed*.

5.3 Surfaces as Manifolds

- A *Non-singular surface* M of dimension k in n -dim Euclidean space is given by a set of $n - k$ equations:

$$f_i(x^1, \dots, x^n) = 0, \quad i = 1, \dots, n - k$$

where $\forall x$ the matrix $\left(\frac{\partial f_i}{\partial x^\alpha} \right)$ has rank $n - k$.

5.4 Orientation of surfaces

5.4.1 Orientation class

- Consider a frame $\tau_1 = (e_1^{(1)}, \dots, e_n^{(1)})$ called an ordered basis and another frame $\tau_1 = (e_1^{(2)}, \dots, e_n^{(2)})$ then we say that they lie in the *same orientation class* if $\det A > 0$ and the *opposite orientation*

class if $\det A < 0$. Where A is defined as:

$$A : e_k^{(1)} \rightarrow e_k^{(2)}$$

5.4.2 Orientability

- A manifold is said to be *orientable* if it is possible to choose at every point of it a single orientation class depending continuously on the points.

A particular choice of such an orientation class for each point is called an orientation of the manifold, and a manifold equipped with a particular orientation is said to be *oriented*.

If no orientation exists the manifold is said to be *non-orientable*

5.5 Two-sided hyper-surface

- A connected $(n - 1)$ -dim sub-manifold of \mathbb{R}^n is called two sided if a single valued continuous field of unit normals can be defined on it.

such a sub-manifold is called a *two-sided hyper-surface*.

6 Lie Groups

6.1 Group

- A *group* is a non-empty set G on which there is defined a binary operation $(a, b) \rightarrow ab$ satisfying the following properties:
 - Closure: If a and b belong to G , then $ab \in G$.
 - Associativity: $\forall a, b, c \in G, \quad a(bc) = (ab)c$.
 - Identity: \exists an element $1 \in G$ st: $a1 = 1a = a, \quad \forall a \in G$
 - Inverse: If $a \in G$ then $\exists a^{-1} \in G$ st: $aa^{-1} = a^{-1}a = 1$.

6.2 Lie Group

- A manifold G is called a *Lie Group* if it has given on it a group operation with the properties that the maps $\varphi : G \rightarrow G$, defined by $\varphi(g) = g^{-1}$ and $\psi : G \times G \rightarrow G$ defined by $\psi(g, h) = gh$, are smooth maps.

6.3 Example of Lie groups

6.3.1 General Linear group

- This is $GL(n, \mathbb{R})$ consisting of all $n \times n$ real matrices with non zero determinant in a region \mathbb{R}^{n^2} .
 $\dim GL(n, \mathbb{R}) = n^2$.

6.3.2 Special Linear group

- This is $SL(n, \mathbb{R})$ consisting of all $n \times n$ real matrices with determinant equal to 1. It is a hyper-surface in \mathbb{R}^{n^2} .

$$\det A = 1, \quad A \in Mat(n, \mathbb{R})$$

$$\dim SL(n, \mathbb{R}) = n^2 - 1.$$

6.3.3 Orthogonal group

- This is $O(n, \mathbb{R})$ consisting of all $n \times n$ real matrices Satisfying:

$$A^T \cdot A = \mathbb{I}, \quad A \in Mat(n, \mathbb{R})$$

$$\dim O(n, \mathbb{R}) = \frac{1}{2}n(n-1).$$

6.3.4 Special Orthogonal group

- This is $SO(n, \mathbb{R})$ consisting of all $n \times n$ real matrices Satisfying:

$$A^T \cdot A = \mathbb{I}, \quad \det(A) = 1, \quad A \in Mat(n, \mathbb{R})$$

$$\dim SO(n, \mathbb{R}) = \frac{1}{2}n(n-1).$$

6.3.5 Pseudo Orthogonal group

- This is $O(p, q, n)$ consisting of all $n \times n$ real matrices Satisfying:

$$A^T \cdot \eta \cdot A = \eta, \quad \det(A) = 1, \quad \eta = \text{diag}\{\underbrace{1, \dots, 1}_p, \underbrace{-1, \dots, -1}_q\}$$

$$\dim O(p, q, n) = \frac{1}{2}n(n-1).$$

6.3.6 Unitary group

- This is $U(n)$ consisting of all $n \times n$ complex matrices Satisfying:

$$A^\dagger \cdot A = \mathbb{I}, \quad A \in \text{Mat}(n, \mathbb{C})$$

$$\dim U(n) = n^2.$$

6.3.7 Special Unitary group

- This is $SU(n)$ consisting of all $n \times n$ complex matrices Satisfying:

$$A^\dagger \cdot A = \mathbb{I}, \quad \det(A) = 1, \quad A \in \text{Mat}(n, \mathbb{C})$$

$$\dim U(n) = n^2 - 1.$$

7 Projective spaces

7.1 Real projective space

- The *real Projective space* $\mathbb{R}P^n$ is the set of all straight lines in \mathbb{R}^{n+1} passing through the origin. Equivalently it is the set of equivalence classes of non-zero vectors in \mathbb{R}^{n+1} where two non-zero vectors are equivalent if they are scalar multiples of one another.
- We may think of $\mathbb{R}P^n$ as obtained from S^n by gluing, that is identifying diametrically opposite points. This means we have the isomorphism $\mathbb{R}P^n \simeq S^n/Z_2$.

7.2 Quaternions

- The set \mathbb{H} of *Quaternions* consists of all linear combinations:

$$q \in \mathbb{H}, \quad q = a\mathbf{1} + b\mathbf{i} + c\mathbf{j} + d\mathbf{k}, \quad a, b, c, d \in \mathbb{R}$$

Where $\mathbf{1}, \mathbf{i}, \mathbf{j}, \mathbf{k}$ are linearly independent. Where these bases satisfy the following multiplications:

$$\begin{aligned} \mathbf{i} \cdot \mathbf{j} &= \mathbf{k} = -\mathbf{j} \cdot \mathbf{i}, & \mathbf{j} \cdot \mathbf{k} &= \mathbf{i} = -\mathbf{k} \cdot \mathbf{j}, & \mathbf{k} \cdot \mathbf{i} &= \mathbf{j} = -\mathbf{i} \cdot \mathbf{k}, \\ \mathbf{i} \cdot \mathbf{i} &\equiv \mathbf{i}^2 = -1, & \mathbf{j} \cdot \mathbf{j} &\equiv \mathbf{j}^2 = -1, & \mathbf{k} \cdot \mathbf{k} &\equiv \mathbf{k}^2 = -1, \\ \mathbf{i} \cdot \mathbf{1} &= \mathbf{i} = \mathbf{1} \cdot \mathbf{i}, & \mathbf{j} \cdot \mathbf{1} &= \mathbf{j} = \mathbf{1} \cdot \mathbf{j}, & \mathbf{k} \cdot \mathbf{1} &= \mathbf{k} = \mathbf{1} \cdot \mathbf{k}, & \mathbf{1} \cdot \mathbf{1} &= \mathbf{1}. \end{aligned}$$

This makes \mathbb{H} an associative algebra over the field of real numbers.

7.3 Complex Projective spaces

- The *complex projective space* $\mathbb{C}P^{\kappa}$ is the set of equivalence classes of non-zero vectors in $\mathbb{C}^{\kappa+\mathbb{K}}$ where two nonzero vectors are equivalent if they are scalar multiples of one another.
- In a similar manner to the real projective space we can identify the isomorphism: $\mathbb{C} \simeq S^{2n+1}/U(1)$.

8 Lie Algebras

8.1 Neighborhood of identity element

- Let G be a Lie group. let the point $g_0 \equiv 1 \in G$ be the identity element of G , and let $T = T_{(1)}$ be the tangent space at the identity element. We can now express the group operations on G in a chart U_0 containing g_0 in terms of local co-ords. We choose co-ords in U_0 so that the identity element is the origin. $g_0 \equiv 1 = (0, \dots, 0)$. then if we let:

$$g_1 = (x^1, \dots, x^n), \quad g_2 = (y^1, \dots, y^n), \quad g_3 = (z^1, \dots, z^n)$$

Which allows us to define the product of two elements:

$$g_1 g_2 = (\psi^1(x, y), \dots, \psi^n(x, y)) = (\psi^i(x, y)) \in U_0$$

An inverse as:

$$g_1^{-1} = (\varphi^1(x), \dots, \varphi^n(x)) = (\varphi^i(x)) \in U_0$$

These functions $\varphi(x), \psi(x)$ satisfy:

$$\begin{aligned} \psi^i(x, 0) &= \psi^i(0, x) = x^i \\ \psi^i(x, \varphi(x)) &= 0 \\ \psi^i(x, \psi(y, z)) &= \psi^i(\psi(x, y), z) \end{aligned}$$

8.1.1 Taylor expansion

- Let $\psi^i(x, y)$ be sufficiently smooth and for $x, y, z \sim \epsilon$:

$$\begin{aligned} \psi^i(x, y) &= x^i + y^i + b_{jk}^i x^j y^k + \mathcal{O}(\epsilon^3) \\ b_{jk}^i &= \left. \frac{\partial^2 \psi^i}{\partial x^j \partial y^k} \right|_{x=y=0} \end{aligned}$$

8.2 Commutator

- Let $\xi, \eta \in T$, and their components in terms of x^i are ξ^i and η^i . Then we can define the *commutator* $[\xi, \eta] \in T$ is defined by:

$$[\xi, \eta]^i = c_{jk}^i \xi^j \eta^k, \quad c_{jk}^i \equiv b_{jk}^i - b_{kj}^i$$

- It has three basic quantities:

- It is *bi-linear* operation on the n -dim vector space T .
- Skew-symmetry: $[\xi, \eta] = -[\eta, \xi]$.
- Jacoby identity: $[[\xi, \eta], \zeta] + [[\zeta, \xi], \eta] + [[\eta, \zeta], \xi] = 0$

8.3 Lie Algebra

- A *Lie algebra* is a vector space \mathcal{G} over a field F with a bi-linear operation $[\cdot, \cdot] : \mathcal{G} \times \mathcal{G} \rightarrow \mathcal{G}$ which is called a commutator or a lie bracket, such that the three axioms above are satisfied.
- This means we can identify the tangent space of a Lie Group at the identity is with respect to the commutator operation of a Lie algebra called the *Lie algebra of the Lie group* G .
- If we choose $\xi = e_j, \eta = e_k$, then combined with the fact that $(e_m) = \delta_m^n$, then we have:

$$[e_j, e_k]^i = c_{jk}^i e_i$$

8.3.1 Structure Constants

- The constants c_{jk}^i which determine the commutation operation on a Lie algebra, and which are skew-symmetric in j, k are called the *structure constants* of the Lie algebra.

9 One parameter subgroups

- A *One parameter subgroup* of a lie group G is defined to be a parametric curve $F(t)$ on the manifold G such that:

$$F(0) = 1, \quad F(t_1 + t_2) = F(t_1)F(t_2), \quad F(-t) = F^{-1}(t)$$

The velocity vector at $F(t)$ is:

$$\frac{dF}{dt} = \frac{dF(t + \epsilon)}{dt} \Big|_{\epsilon=0} = \frac{d}{d\epsilon}(F(t)F(\epsilon)) \Big|_{\epsilon=0} = F(t) \frac{dF(\epsilon)}{d\epsilon} \Big|_{\epsilon=0}$$

Hence:

$$\dot{F}(t) = F(t)\dot{F}(0) \quad \text{or} \quad F^{-1}(t)\dot{F}(t) = \dot{F}(0)$$

i.e. the induced action of left multiplication by $F^{-1}(t)$ sends $\dot{F}(t)$ to $\dot{F}(0) = \text{const} \in T$.

- conversely, $\forall A \in T$ the equation $F^{-1}(t)\dot{F}(t) = A$ is satisfied by a unique one-parameter subgroup $F(t)$ of G . If G is a matrix group then $F(t) = \exp(At)$.

9.1 Co-ords of the first kind

- One parameter subgroups can be used to define so called *canonical* in a neighborhood of the identity of a Lie group G .
- Let A_1, \dots, A_n form a basis for the Lie algebra T . Then $\forall A = \sum_i A_i x^i \in T \exists$ a one parameter group $F(t) = \exp(At)$. To the point $F(1) = \exp(A)$ we assign as co-ords co-officiants x^1, \dots, x^n , which gives us a system of co-ords in a sufficiently small neighborhood of $g_0 = 1 \in G$. These are called the *canonical co-ords of the first kind*.

9.2 Co-ords of the second kind

- Another system of co-ords is obtained by introducing $F_i(t) = \exp(A_i t)$ and representing a point g sufficiently close to g_0 as:

$$g = F_1(t_1)F_2(t_2) \cdots F_n(t_n)$$

for small t_1, \dots, t_n . Assigning co-ords $x^1 = t_1, \dots, x^n = t_n$ to the point g , we get the *canonical co-ords of the second kind*.

10 Linear Representations

10.1 Representations

- A *Linear representation* of a group G of $\dim G = n$ is a homomorphism:

$$\rho : G \rightarrow GL(r, \mathbb{R}), \quad \text{or} \quad \rho : G \rightarrow GL(r, \mathbb{C})$$

- Given a representation ρ of G the map:

$$\chi_\rho : G \rightarrow \mathbb{R}, \quad \text{or} \quad G \rightarrow \mathbb{C}$$

defined by:

$$\chi_\rho(g) = \text{tr}(\rho(g))$$

is called the *character* of the representation ρ .

- A representation ρ of G is said to be *irreducible* if the vector space \mathbb{R}^r contains no proper subspace invariant under the matrix group $\rho(G)$.

10.1.1 Matrix Invariance

- A subspace W of the representation space \mathbb{R}^r is called *invariant under the matrix group* $\rho(G)$ (or simply G invariant) if:

$$\rho(G)W \subset W, \quad \forall g \in G$$

Then we can restrict ρ to W and get a *subrepresentation*.

10.2 Schur's Lemma

- Let $\rho_i : G \rightarrow GL(r_i, \mathbb{R})$, $i = 1, 2$ be two irreducible representations (irreps) of a group G . If $A : \mathbb{R}^{r_1} \rightarrow \mathbb{R}^{r_2}$ is a linear transformation changing ρ_1 to ρ_2 , i.e. stratifying:

$$A\rho_1(g) = \rho_2(g)A, \quad \forall g \in G$$

Then either A is the zero transformation or else a bijection, in which case $r_1 = r_2$.

10.3 Push Forward Representation

- If G is a Lie group and a representation $\rho : G \rightarrow GL(r, \mathbb{R})$ is a smooth map, then the push-forward map ρ_* is a linear map from the Lie algebra $\mathfrak{g} = T_{(1)}$ to the space of all $r \times r$ matrices:

$$\rho_* : \mathfrak{g} \rightarrow \text{Mat}(r, \mathbb{R})$$

It can then be shown that this means ρ_* is a *representation* of the Lie algebra \mathfrak{g} , i.e. that it is a Lie algebra homomorphism. Meaning it is linear and preserves the commutators $\rho_*[\xi, \eta] = [\rho_*\xi, \rho_*\eta]$.

10.4 Faithful

- A representation $\rho : G \rightarrow GL(r, \mathbb{R})$ is called *faithful* if it is one to one i.e. if its Kernel is trivial. So $\rho(g) \neq \mathbb{I}$ unless $g = g_0$.
- If a Lie group has a faithful representation then it can be realized as a matrix Lie group.

10.5 Inner automorphism

- For each $h \in G$ the transformation $G \rightarrow G$ defined by $g \rightarrow hgh^{-1}$ is called the *inner automorphism* of G determined by h .
- Any inner automorphism does not move the identity element. i.e. $g_0 = hg_0h^{-1}$ and therefor the push forward (induced linear) map of the tangent space T to G at g_0 is a linear transformation of T denoted by:

$$Ad_h : T \rightarrow T$$

it satisfies the following:

- $Ad_{g_0} = id$, where id is the identity transformation of T .
- $Ad_{h_1}Ad_{h_2} = Ad_{h_1h_2}$ for all $h_1, h_2 \in G$. because $h_1h_2gh_2^{-1}h_1^{-1} = (h_1h_2)g(h_1h_2)^{-1}$.
- Choosing $h_1 = h, h_2 = h^{-1}$, we get that $Ad_{h^{-1}} = Ad_h^{-1}$
- This means that the map $h \rightarrow Ad_h$ is a *linear representation* of the group G . i.e. a homomorphism to a group of linear transformations, $Ad : G \rightarrow GL(n, \mathbb{R}), h \rightarrow Ad_h = Ad(h)$. This representation of G is called *Adjoint*.

10.6 One Parameter Adjoint

- Let $F(t) = e^{At}$ be a one parameter subgroup of a Lie group G . Then $Ad_{F(t)}$ is a one parameter subgroup of $GL(n, \mathbb{R})$.

The vector $\left. \frac{d}{dt} Ad_{F(t)} \right|_{t=0}$ lies in the Lie algebra $\mathfrak{g} \sim Mat(n, \mathbb{R})$ of the Group $GL(n, \mathbb{R})$ and can be regarded as a linear operator.

- This operator is denoted ad_A and is given by:

$$ad_A : \mathbb{R} \rightarrow \mathbb{R}, \quad B \mapsto [A, B], \quad B \in T \simeq \mathbb{R}^n$$

11 Simple Lie Algebras and Forms

11.1 Simple & Semi-Simple

- A Lie algebra $\mathfrak{g} = \{\mathbb{R}^n, c_{jk}^i\}$ is said to be *simple* if it is *non-commutative* and has *no proper ideals*, i.e. subspaces $\mathcal{I} \neq \mathfrak{g}, 0$ for which $[\mathcal{I}, \mathfrak{g}] \subset \mathcal{I}$.
- It is instead called *semi-simple* if we can write $\mathfrak{g} = \mathcal{I}_1 \otimes \mathcal{I}_2 \otimes \cdots \otimes \mathcal{I}_k$ Where the \mathcal{I}_j are ideals which are simple as Lie algebras. These ideals are pairwise commuting $[\mathcal{I}_i, \mathcal{I}_j] = 0, \quad i \neq j$.

A Lie group is defined to be simple or semi-simple according to its Lie algebra.

- A theorem that can be proven is that if the Lie algebra \mathfrak{g} of a Lie group G is simple, then the linear representation $Ad : G \rightarrow GL(n, \mathbb{R})$ is *irreducible*, i.e. \mathfrak{g} has no proper invariant sub-spaces under the group of inner automorphisms Ad_G .

11.2 Killing Form

- The *Killing form* on an arbitrary Lie algebra \mathfrak{g} is defined (up to a sign) by:

$$\langle A, B \rangle = -\text{tr}(ad_A ad_B)$$

- If the Killing form of a Lie algebra is positive definite then the Lie algebra is semi-simple.
- We also have that a Lie algebra is semi-simple if and only if its Killing form is non-degenerate.

12 Group Actions

12.1 Left and Right actions

- We say that a Lie group G is represented as a *group of transformations* of a manifold M , or has a *left action* on M if:
 - There is associated with each of its elements g a diffeomorphism from M to itself. $x \mapsto \mathcal{T}_g(x)$, $x \in M$. Such that $\mathcal{T}_g\mathcal{T}_h = \mathcal{T}_{gh}$, $\forall g, h \in G$.
 - $\mathcal{T}_g(x)$ depends smoothly on the arguments g, x i.e. the map $(g, x) \mapsto \mathcal{T}_g(x)$ is a smooth map from $G \times M \rightarrow M$.
- The Lie group is said to have *Right action* on M if the above definition is valid with $\mathcal{T}_g\mathcal{T}_h = \mathcal{T}_{hg}$.

12.2 Transitivity

- The action of a group G on M is said to be *transitive* if for every two points $x, y \in M$ there exists an element of G such that $\mathcal{T}_g(x) = y$.

To show that an action of a group on a manifold is transitive it is sufficient to choose any point of M as a reference point x_0 , and to prove that for any point $y \in M$ there exists an element $g \in G$ such that $y = \mathcal{T}_g(x_0)$.

12.2.1 Homogeneity

- A manifold on which a Lie group acts transitively is called a *homogeneous space* of the Lie group.
- In particular, G is a homogeneous space for itself, e.g. as $h \rightarrow \mathcal{T}_g(h) = gh$, $h \in G$. G is called the *principle* homogeneous space.

12.2.2 Isotropy group

- Let x be any point of a homogeneous space M of a Lie group G . The *isotropy group* (or *stationary group*) H_x of the point x is the stabilizer of x under the action of G :

$$H_x = \{h | \mathcal{T}_h(x) = x\}$$

- All isotropy groups H_x of points x of a homogeneous space are isomorphic.
- There is a one to one correspondence between the points of a homogeneous space M of a group G , and the left cosets gH of H in G , where H is the isotropy group and G acts on the left. Thus we can write $M \simeq G/H$, i.e. M is a diffeomorphic to the quotient space G/H .

12.3 Examples of Homogeneous spaces

12.3.1 Stiefel manifolds

- For each n, k the Stiefel manifold $V_{n,k}$ has as its points *all orthonormal k -frames* $x = (e_1, \dots, e_k)$ of k vectors e_a in \mathbb{R}^n .
- The dimension of $V_{n,k}$ is $nk - \frac{1}{2}k(k+1)$ and $V_{n,k} \simeq O(n)/O(n-k) \simeq SO(n)/SO(n-k)$.

12.3.2 Real Grassmanian manifolds

- The points of $G_{n,k}$ are the k dimensional planes passing through the origin of \mathbb{R}^n .
- It can be shown that $G_{n,k} \simeq O(n)/(O(k) \times O(n-k)) \simeq G_{n,n-k}$. The dimension of $G_{n,k}$ is $(n-k)k$

13 Vector Bundles

13.1 Tangent Bundle

- The *tangent bundle* $T(M)$ of an n dimensional manifold M is a $2n$ dimensional manifold defined as follows:
 - The points of $T(M)$ are the pairs $(x, \xi), x \in M, \xi \in T_x M$.
 - Given a chart U_q of M with the local co-ords (x_q^i) , the corresponding chart U_q^T of $T(M)$ is the set of all pairs (x, ξ) where:

$$x = (x_1^1, \dots, x_q^n) \in U_q, \quad \xi = \xi_q^i \frac{\partial}{\partial x_q^i} \in T_x M$$

$$\text{with local co-ords } (y_q^1, \dots, y_q^{2n}) = (x_q^1, \dots, x_q^n, \xi_q^1, \dots, \xi_q^n) = (x_q^i, \xi_q^i).$$

- This tangent bundle is a smooth oriented manifold.

13.2 Cotangent Bundle

- The *cotangent bundle* $T^*(M)$ of an n dimensional manifold M is a $2n$ dim manifold defined as follows:
 - The points $T^*(M)$ are the pairs $(x, p), x \in M$ and p a co-vector at the point x , so $p \in T_x^* M$.
 - Given a chart U_q of M with the local co-ords (x_q^i) , the corresponding chart $U_x^{T^*}$ of $T^* M$ is the set of all pairs (x, p) , where:

$$x = (x_1^1, \dots, x_q^n) \in U_q, \quad p = p_{qi} dx_q^i \in T_x^* M$$

$$\text{with local co-ords } (y_q^1, \dots, y_q^{2n}) = (x_q^1, \dots, x_q^n, p_{q1}, \dots, p_{qn}) = (x_q^i, p_{qi}).$$

- This cotangent bundle is a smooth oriented manifold.

13.3 Symplectic Manifold

- The existence of a metric on M gives rise to a map:

$$T(M) \rightarrow T^*(M) : (x^i, \xi^i) \mapsto (x^i, g_{ij} \xi^i)$$

- Since $\omega = p_i dx^i$, a differential one-form on M , is invariant under a change of co-ords of $T^*(M)$, it is a differential form on $T^*(M)$.

Its differential $\Omega = d\omega = dp_i \wedge dx^i$ is a *non-degenerate closed*, ($d\Omega = 0$), 2-form on $T^*(M)$.

- Thus $T^*(M)$ is a *symplectic* manifold, i.e. it is equipped with a closed non-degenerate 2-form.

14 Vector and Tensor Fields

14.1 Vector Field

- A *vector field* is a map that specifies a unique vector at each point x of the manifold M :

$$\xi : M \rightarrow T(M), \quad x \mapsto \xi_x \in T_x M$$

A vector field intersects each tangent space of $T(M)$ at one and only one point, i.e. a vector field is a curve which is no-where parallel to a tangent space. It is a *cross section* of $T(M)$.

- A vector field can be understood as a differential operator that maps a scalar function to a scalar function on M :

$$\xi(f) = \xi^i \frac{\partial f}{\partial x^i}.$$

- These maps are linear and satisfy the Leibniz rule. This means they are *derivations*.

14.2 Tensor Field

- A *Tensor field* of type (r, s) assigns a unique tensor of type (r, s) to each point x of the manifold M :

$${}^{(r,s)}\xi : M \rightarrow T^{(r,s)}(M), \quad x \mapsto {}^{(r,s)}\xi_x \in T_x^{(r,s)} M$$

. It is a cross section of $T_x^{(r,s)} M$.

14.3 Commutator or Lie Bracket

- Consider the composition $\xi(\eta(f)) = \xi^i \frac{\partial}{\partial x^i} \left(\eta^j \frac{\partial f}{\partial x^j} \right) = \xi^i \frac{\partial \eta^j}{\partial x^i} \frac{\partial f}{\partial x^j} + \xi^i \eta^j \frac{\partial^2 f}{\partial x^i \partial x^j}$. The second term makes this not a vector field.
- This motivates us to define the *commutator* or *Lie Bracket* define by:

$$[\xi, \eta](f) \equiv \xi(\eta(f)) - \eta(\xi(f)) = \left(\xi^i \frac{\partial \eta^j}{\partial x^i} - \eta^i \frac{\partial \xi^j}{\partial x^i} \right) \frac{\partial f}{\partial x^j}$$

This is a vector field that satisfies:

$$\begin{aligned} [\xi, \eta] &= -[\eta, \xi] \\ [\xi, \eta + \zeta] &= [\xi, \eta] + [\xi, \zeta] \\ [\xi, f\eta] &= f[\xi, \eta] + \xi(f)\eta \\ [\xi, [\eta, \zeta]] + [\eta, [\zeta, \xi]] + [\zeta, [\xi, \eta]] &= 0 \end{aligned}$$

Thus the vector space equipped with the commutator operation is an *infinite dimensional Lie algebra*.

14.4 Integral curves

- let $\xi^i(x)$ be a vector field on M . Consider the autonomous (meaning the equations have no explicit dependence on t) system of differential equations:

$$\dot{x}^i \equiv \frac{dx^i}{dt} = \xi^i(x^1(t), \dots, x^n(t)), \quad i = 1, \dots, n$$

- The solutions $x^i(t)$ to this system are called the *integral curves* of the vector field ξ^i . The vector field $\xi^i(x)$ is comprised of tangent vector to the integral curves.

14.4.1 Flows and Velocity Fields

- A local abelian one parameter subgroup of diffeomorphisms F_t is called the *flow* generated by the vector field ξ^i . Where $F_t^i(x_0^1, \dots, x_0^n) = x^i(t, x_0^1, \dots, x_0^n)$.
- This can be reversed to say that given a one parameter local group of diffeomorphisms we can define its *velocity fields* to be the vector field:

$$\xi^i = \left(\frac{d}{dt} F_t^i \right)$$

- Note that in general two flows do not commute and in fact the commutator measures the discrepancy between the points obtained by following the integral curves of two different vector fields in different orders.
- The vectors comprising a co-ord induced basis commute because all the partial derivatives do, The converse is also true, if all the elements of a basis for vector fields commute then the basis is co-ordinate induced.

14.5 Exponential function of Vector Fields

- A one parameter subgroup of diffeomorphisms $F_t(x)$ with associated vector field $\xi(x)$ is defined to *act on smooth functions* $f = f(x)$ as follows:

$$(F_t f)(x) = f(F_t(x)).$$

- The *exponential function* of a vector field ξ is the operator:

$$\exp(t\partial_\xi) = 1 + \partial_\xi + \frac{t^2}{2}(\partial_\xi^2) + \dots$$

Where ∂_ξ is the directional derivation operator in the direction of ξ .

- The action of $e^{t\partial_\xi}$ on functions $f(x)$ is defined as:

$$\exp(t\partial_\xi)f = f + \partial_\xi f + \frac{t^2}{2}(\partial_\xi^2)f + \dots$$

‘ $\forall t$ for which this series converges.

- For analytic vector fields $\xi(x)$ and analytic functions $f(x)$ the exponential function of $\xi(x)$ i.e. $e^{t\partial_\xi}$, coincides for sufficiently small t with the action of F_t on f :

$$e^{\partial_\xi} f = f(F_t(x))$$

15 The Lie Derivative

- The motivation for this is that we want to define the action of a flow F_t generated by ξ on Tensors. The problem with this is that we have no way of comparing tensors at *different points* on the manifold. To fix this we switch to active transformations where the points x^i stay fixed and we instead change the co-ordinate system by acting on the basis with $F_t^{-1} = F_{-t}$.

15.1 Action of Flows on Tensors

- A one parameter subgroup of diffeomorphisms $F_t(x)$ with associated vector field $\xi(x)$ is defined to *act on smooth tensors* $T = (T_{j_1, \dots, j_q}^{i_1, \dots, i_p})$ of type (p, q) as follows:

$$(F_t T)_{j_1, \dots, j_q}^{i_1, \dots, i_p}(x) = T_{l_1, \dots, l_q}^{k_1, \dots, k_p}(y) \frac{\partial y^{l_1}}{\partial x^{j_1}} \cdots \frac{\partial y^{l_q}}{\partial x^{j_q}} \frac{\partial x^{i_1}}{\partial y^{k_1}} \cdots \frac{\partial x^{i_p}}{\partial y^{k_p}}$$

Where $y^i = F_t^i(x)$.

15.2 Lie Derivative

- The Lie derivative of a tensor $T = (T_{j_1, \dots, j_q}^{i_1, \dots, i_p})$ along a vector field ξ is the tensor $\mathcal{L}_\xi T$ given by:

$$\mathcal{L}_\xi T_{j_1, \dots, j_q}^{i_1, \dots, i_p} = \left[\frac{d}{dt} (F_t T)_{j_1, \dots, j_q}^{i_1, \dots, i_p} \right]_{t=0}$$

15.2.1 Killing Vector

- If a vector ξ satisfies $\mathcal{L}_\xi g_{ij} = 0$, where g_{ij} is the metric, then ξ is called a *Killing Vector*.
- A lemma of this definition is that the Killing vector fields of a (pseudo-)Riemann manifold form a Lie algebra with respect to the Lie bracket given by the commutator of the two fields.

$$\mathcal{L}_{[\xi, \eta]} g_{ij} = [\mathcal{L}_\xi, \mathcal{L}_\eta] g_{ij} = 0$$

16 Covariant Differentiation

- We need to define a different type of differentiation as π_μ

